

Research Article

Phytochemical analysis of red alga *Champia parvula* (C. Agardh) collected from Mandapam coast of Tamil Nadu, India

Vinoth Kumar, R, Murugesan, S* and Bhuvaneshwari, S

Unit of Algal Biotechnology and Bionano Technology, PG and Research Dept of Botany, Pachaiyappa's College, Chennai – 600 030.

***Correspondence Info:**

Murugesan, S
Unit of Algal Biotechnology and
Bionano Technology, PG and Research
Dept of Botany, Pachaiyappa's College,
Chennai – 600 030.
Email: smurugesan5@gmail.com

Keywords:

Phytochemical analysis,
Champia parvula,
GC-MS, fatty acid.

Abstract

The marine red alga *Champia parvula* showed the phytochemical constituents like sterols, glycosides, anthroquinones, phenols, alkaloids, triterpenoids, tannins, saponins, flavonoids, steroids. Flavonoid compounds have rutin, quercetin, kamferol and phenol compounds have gallic acid and cinnamic acid. Similarly, the fatty acids have palmitic acid, margaric acid, stearic acid, oleic acid, linolenic acid, alpha linolenic acid, morotic acid were also present. Among the phytochemical contents the triterpenoids and glycosides are present in high. Among the seven fatty acid, stearic acid ($6.03 \pm 0.012\%$) and morotic acid ($5.58 \pm 0.004\%$) were identified. The aim of the present study is to evaluate the phytochemical constituents of the marine red alga *Champia parvula*.

1. Introduction

The creature of nature possesses a magnificent primitive organism such as seaweeds or marine macroscopic algae which form an important component of marine vast living organisms. In several Asian countries seaweed consumption as vegetables in human diets has been a common practice [1]. They are well known as an excellent source of biologically active compounds. People who are living in coastal areas consume fresh and dry seaweeds [2]. Algae are very closely associated with the human health are being exhaustively used in numerous ways as a source of food, feed, fertilizer, medicine and chiefly for economically important phycocolloids [3,4]. The phytochemicals from marine algae are extensively used in various industries such as food, confectionery, textile, pharmaceutical, dairy and paper industry for mostly used as gelling, stabilizing and thickening agents [5]. The bioactive compounds isolated from seaweeds are used as medicine as well as food all over the world since 13,000 years [6]. Seaweeds are rich in antioxidant such as carotenoids, pigments, polyphenols, enzymes and diverse functional polysaccharide [7], Seaweeds have some of valuable medicinal compounds such as antibiotics [8] laxative, anticoagulants, [9]. Antiulcer products and suspending agents in radiological preparation [10]. Algae are the source of amino acids, terpenoids, phlorotannins, steroids, phenolic compounds, halogenated ketones and alkanes, cyclic polysulphides, fatty acids, acrylic acid [11,12] proteins, polysaccharides, vitamins, minerals and fibers [13,14]. Marine macroalgae are considered as an excellent source of bioactive compounds which has a broad range of biological activities including antibacterial [15-23], antiviral [24], antifungal [25], anticoagulant [26] antitumor [27] and anti-inflammatory activities [28]. Alginate, carrageenan and agar obtained from seaweeds have been used in medicine and pharmacy [29]. The presence of these secondary metabolites in seaweeds is highly evident of their pharmaceutical potential. Thus, the present investigation was carried to evaluate the phytochemical constituents of marine red alga *Champia parvula*.

2. Material and Methods

2.1 Sample collection and extraction methods

The marine red alga *Champia parvula* collected from the intertidal region of the Mandapam coastal water and immediately brought to the laboratory in plastic bags containing water to prevent evaporation. Then the sample was washed thoroughly with sea water to remove extraneous materials. Samples were dried at 37°C and ground in an electric mixer. The coarse powder was soaked in the methanol for 24 hrs after that the mixture was filtered by using filter paper to separate the extract. The extract was then concentrated and dried by a rotary evaporator. The quantitative phytochemical analyses of the sample were carried out by standard procedure.

2.2 Quantitative phytochemical analysis

The amount of phytochemicals present in the red alga *Champia parvula* extract was quantitatively determined by standard procedure.

2.3 Determination of total flavonoid compounds

This method is based on the formation of the flavonoids - aluminium complex, which has an absorptive maximum at 415 nm. The alga extract of 100 µL in methanol (10 mg/mL) was mixed with 100 µL of 20 % aluminum trichloride in methanol and a drop of acetic acid, and then diluted with methanol to 5 mL. The absorption at 415 nm was read after 40 minutes. Blank samples were prepared from 100 mL of alga extract and a drop of acetic acid, and then diluted to 5 mL with methanol.

2.4 Determination of total phenolic compounds

The extract, 100 mg of the sample was weighed accurately and dissolved in 100 mL of double distilled Water (DDW). 1 mL of this solution was transferred to a test tube, then 0.5 mL 2N of the Folin-Ciocalteu reagent and 1.5 mL 20% of Na₂CO₃ solution was added and ultimately the volume was made up to 8 mL with (DDW) followed by vigorous shaking and finally allowed to stand for 2 hours after which the absorbance was taken at 765 nm [30].

2.5 Determination of Fatty acids compounds

The 100 mg of the extract of the sample was weighed accurately and dissolved in 100 mL of Double Distilled Water (DDW) 1 mL of this solution was transferred to a test tube, then 1 mL of ether solution was added and ultimately the volume was made up to 8 mL with (DDW) followed by vigorous shaking and finally allowed to stand for 2 hrs after which the absorbance was taken at 765 nm [31]. These data were used to estimate the total fatty acid content.

2.6 Statistical analysis

All the data for phytochemical analysis subjected to analysis of variance (ANOVA) using SPSS version (16.0). The results were expressed as mean ± standard error (SE).

3. Results and Discussion

There are many studies were reported on the presence of different phytochemical compounds of marine red algae collected from the coastal area [16,32]. The present investigation reveals that the red alga *Champia parvula* extract contained 20.17 mg of sterols, 35.33 mg of glycosides, 10.43 mg of anthroquinones, 25.50 mg of phenols, 12.56 mg of alkaloids, 55.33 mg of triterpenoids, 14.20 mg of tannins, 22.50 mg of saponins, 10.17 mg of flavonoids, 24.30 mg of steroids, and 5.45 mg of fatty acids. The flavonoid contents were 5 mg of rutin, 2.87 mg of quercetin, and 2.10 mg of kamferol and phenol contents were 3.40 mg of gallic acid and 2.02 mg of cinnamic acid and fatty acid content are 1.82 mg of palmitic acid, 2.07 mg of margaric acid, 6.03 mg of stearic acid, 1.16 mg of oleic acid, 3.86 mg of linolenic acid, 4.89 mg of alpha-linolenic acid, and 5.58 mg of moroctic acid. The extract showed different amount of phytochemicals. Similarly phytochemical analysis shows triterpenoids are rich than other phytochemicals. (Table 1). The composition of flavonoids (Table 2), phenols (Table 3) and fatty acids (Table 4) were rich in the red alga *Champia parvula* compared to other bioactive compounds.

Table 1: Quantitative analysis of phytochemicals composition (mg/g) in *Champia parvula*

S. No.	Phytochemicals (mg/g)	Mean \pm S.E
1	Sterols	20.17 \pm 0.01
2	Glycosides	35.33 \pm 0.14
3	Anthroquinones	10.43 \pm 0.00
4	Phenols	25.50 \pm 0.11
5	Alkaloids	12.56 \pm 0.17
6	Triterpenoids	55.33 \pm 0.14
7	Tannins	14.20 \pm 0.05
8	Saponins	22.50 \pm 0.11
9	Flavonoids	10.17 \pm 0.01
10	Steroids	24.30 \pm 0.11
11	Fattyacids	05.45 \pm 0.01
F – Value		1498.0
P- Value		0.000

Table 2: Quantitative analysis of Flavonoid composition (mg/g) in *Champia parvula*

S. No.	Flavonoid composition (mg/g)	Mean \pm S.E
1	Rutin	5.00 \pm 0.05
2	Quercetin	2.87 \pm 0.06
3	Kamferol	2.10 \pm 0.00
F – Value		101.9
P- Value		0.000

Table 3: Quantitative analysis of phenol composition (mg/g) in *Champia parvula*

S. No.	Phenol composition (mg/g)	Mean \pm S.E
1	Gallic acid	3.40 \pm 0.031
2	Cinnamic acid	2.02 \pm 0.020
F – Value		19.455
P- Value		0.000

Table 4: Quantitative analysis of Fatty acid compositions (mg/g) in *Champia parvula*

S. No.	Fatty acids	Mean \pm S.E
1	Palmitic acid	1.82 \pm 0.064
2	Margaric acid	2.07 \pm 0.014
3	Stearic acid	6.03 \pm 0.012
4	Oleic acid	1.16 \pm 0.017
5	Linolenic acid	3.86 \pm 0.052
6	Alpha linolenic acid	4.89 \pm 0.006
7	Moroctic acid	5.58 \pm 0.004
F- Value		3526.3
P- Value		0.000

Flavonoids are proved to have antitumour and antioxidant properties [33]. Glycosides are known to lower the blood pressure [34]. In the present study the *Champia parvula* showed the presence of glycosides. Tannins bind to proline rich protein and interfere with the protein synthesis [35]. Tannin containing remedies are in use as antihelminthic [36], antioxidant [37], antimicrobial and antiviral [38] and for cancer treatment [39] terpenoids have antioxidant activity [40]. Alkaloids are commonly found to have antimicrobial [41-43], cytotoxic [44] and antiplasmodic properties [45,46]. Steroids isolated from marine algae have medicinal value [10,47] saponins have unique residue like 2, 3-dihydro-2, 5-dihydroxy-6-methyl-4H-pyran-4-one (DDMP), which allows saponins to scavenge superoxides by forming hydroperoxide intermediates which prevent bio-molecular damage [48].

The presence of different bioactive constituents in the methanolic extract of *Champia parvula* could contribute to different biological activities. The phenol compounds were reported to be responsible for the antioxidant activity in

Spyridia fusiformis and *Chondrococcus hornemanni* [49], The presence of glycosides, carbohydrates, tannins, alkaloids, steroids, terpenoids, phytosterols and saponins were recorded in marine alga [50] phenolic compounds are commonly found in plants, including seaweeds, and have been reported to have a wide range of biological activities including antioxidant properties [51]. Phytochemical screening of marine red alga *Champia parvula* is very important to identify new sources of therapeutically and industrially important compounds. It is important to initiate an urgent step for the screening of the red alga for secondary metabolites.

4. Conclusion

In conclusion, the results of this phytochemical investigation suggest that the marine red alga *Champia parvula* contains important phytochemicals like phenols, flavonoids and fatty acids, which may contribute to its biological activities. The pharmacological properties require further investigation of these active ingredients by implementing techniques of extraction, purification, separation, compound isolation and identification.

Acknowledgement

The authors are grateful to the Department of Science and Technology (DST), New Delhi for its financial support (Ref. No: Inspire fellowship IF140803).

References

- [1] Nisizawa, K. Seaweeds Kaiso. Japan Seaweed Association. Publication, Kochi, Japan. 2002; Page pp.23.
- [2] Cox, S., Abu-Ghannam, N. and Gupta, S. An assessment of the antioxidant and antimicrobial activity of six species of edible Irish seaweeds. *International Food Research Journal* 2010; 17: 205-220.
- [3] Levering, T.; Hoppe, H. A and Schmid, O.J. Marine Algae. A survey of research and Utilization. Gramm be Gruyter & Co., Hamburg, 1969; 1-421.
- [4] Chapman, V.J. Seaweeds and their uses. The Camelot Press Ltd., Methuen and Co Ltd., London and Southampton, II Edn.: 1970; 63-85 pp.
- [5] Seenivasan R; Rekha M; Geetha S. Antibacterial Activity and Phytochemical Analysis of Selected Seaweeds from Mandapam Coast, India. *Journal of Applied Pharmaceutical Science*. 2012; 2(10): 159-169.
- [6] Radhika D; Veerabahu C; Priya R. Antibacterial Activity Of Some Selected Seaweeds From The Gulf of Mannar Coast, South India *Asian J. Pharm. Clin. Res.*, 2012; 5(4), 89-90.
- [7] Muhammad Shaiq Ali, Viqar Uddin Ahmad, Farah Mazhar, Iqbal Azhar, Khan Usmanghani. Some Chemical Constituents from Marine Algae of Karachi Coast (Arabian Sea) *Turk J Chem* 1999; 23: 181-183.
- [8] Swaminathan, S., Murugesan, S., Damodarkumar, S., Dhamotharan, R and Bhuvanewari, Synthesis and Characterization of Gold Nanoparticles from Alga *Acanthophora specifera* (VAHL) Boergesen. *International Journal of Nano Science and Nanotechnology*, 2011; 2 (2): 85-94.
- [9] Koyanagi S, Tanigawa N, Nakagawa H, Soeda S and Shimeno H. Oversulfation of fucoidan enhances its anti-angiogenic and antitumor activities. *Biochem Pharmacol* 2003; 65: 173-179.
- [10] Rajasulochana P; Dhamotharan R; P Krishnamoorthy; S Murugesan. Antibacterial Activity of the Extracts of Marine Red and Brown Algae. *Journal of American Science*, 2009; 5(3): 20-25.
- [11] Mtolera, M.S.P. and Semesi, A.K. Antimicrobial activity of extracts from six green algae from Tanzania. In: Current Trends in Marine Botanical Research in the East African Region (eds M. Bjérk, A.K. Semesi, M. Pedersen and B. Bergman). Gotab AB, Uppsala, Sweden, 1996; pp. 211-217.
- [12] Taskin, E., Ozturk, M., Taskin, E. and Kurt, O. Antibacterial Activities of Some Marine Algae from the Aegean Sea (Turkey). *African Journal of Biotechnology*, 2007; 6: 2746-2751.
- [13] Lahaye M. Marine algae as sources of fibres: Determination of soluble and insoluble dietary fibre contents in some 'sea vegetables'. *J. Sci. Food Agric.*, 1999; 54, 587-594.
- [14] Darcy-Vrillon B. Nutritional aspects of the developing use of marine macroalgae for the human food industry. *Int. J. Food Sci. Nutr.* 1993; 44: 523-535.
- [15] Burkholder PR., Burkholder LM., Almodovar LR. Antibiotic activity of some marine algae of Puerto Rico. *Bot. Mar.* 1960; 2: 149- 156.

- [16] Sachithanathan K., Sivapalan A. Antibacterial properties of some marine algae of Sri Lanka. Bulletin of Fisheries Research Station, Sri Lanka. 1975; 26:5-9.
- [17] Mahasneh I., Jamal M., Kashashneh, M., Zibdeh M. Antibiotic activity of marine algae against multiantibiotic resistant bacteria. *Microbios.* 1995; 83:23-26.
- [18] Tuney, I.; Çadirci, B.H.; Unal, D.; Sukatar, A. Antimicrobial activities of the extracts of marine algae from the coast of Urla. *Turk. J. Biol.* 2006; 30: 171–175.
- [19] Karthikaidevi G; Manivannan K; Thirumaran G; Anantharaman P; Balasubramanian T. Antibacterial Properties of Selected Green Seaweeds from Vedalai Coastal Waters; Gulf of Mannar Marine Biosphere Reserve. *Global Journal of Pharmacology*, 2009; 3(2): 107-112.
- [20] Rizvi MA. Comparative Antibacterial Activities of Seaweed Extracts From Karachi Coast, Pakistan. *Pakistan Journal of Pharmacology*, 2010; 27(2), 53-57.
- [21] Lavanya R; Veerapan N. Antibacterial Potential of Six Seaweeds Collected from Gulf of Mannar of Southeast Coast of India. *Advances in Biological Research* 2011; 5(1): 38-44.
- [22] Sumathi S, and Krishnaveni M. Preliminary screening, antioxidant and antimicrobial potential of *Chaetomorpha antennina* and *Caulerpa scalpelliformis* in vitro study. *International Journal of Environmental sciences*, 2012; 2(3): 2312-2320.
- [23] Alghazeer R; Whida F; Abduelrhman E; Gammoudi F; Azwai S. Screening of antibacterial activity in marine green, red and brown macroalgae from the western coast of Libya Natural Science 2013; 5(1): 7-14.
- [24] Richards JT; Kern ER; Glasgow LA; Overall JC; Deign EF; Melvin MT. Antiviral Activity of Extracts from Marine Algae. *Antimicrobial Agents Chemother.* 1978; 14, 24-30.
- [25] Oumaskour K, Boujaber N, Etahiri S; Assobhei O. Screening of antibacterial and antifungal activities in green and brown algae from the coast of Sidi Bouzid (El Jadida, Morocco). *African Journal of Biotechnology*, 2012; 11(104): 16831-16837.
- [26] Yasantha A; Ki-Wan L; Se-Kwon K; You-Jin J. Anticoagulant activity of marine green and brown algae collected from Jeju Island in Korea. *Bioresour. Technol.*, 2007; 98: 1711-1716.
- [27] Abirami RG; Kowsalya S. Anticancer Activity of Methanolic and Aqueous Extract of *Ulva Fasciata* In Albino Mice. *International Journal of Pharmacy and Pharmaceutical Science*, 2012; 4(2): 681-684.
- [28] Boonchum W; Peerapornpisal Y; Kanjanapathi D; Pekkoh J; Amornlerdpison D; Pumas C; Sangpaiboon P; Vacharapiyasophon P. Antimicrobial and anti inflammatory properties of various seaweeds from the gulf of Thailand. *Int. J. Agric. Biol.*, 2011; 13: 100-104.
- [29] Siddhanta AK; Mody KH; Ramavat BK; Chauhan VD; Garg HS; Goel AK; Doss MJ; Srivastava MN; Patnaik GK, Kamboj VP. Bioactivity of marine organisms: Part VIII-Screening of some marine flora of western coast of India. *Indian J. Exp. Biol.*, 1997; 36, 638-643.
- [30] Taga, M.S. Miller, E.E and Pratt D. F. *Journal of the American Oil Chemists Association.* 1984; 61: 928-931.
- [31] Zhishen, J.; Mengheng, T and Jianming, W. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry.* 1999; 64: 555-559.
- [32] Vidhyavathi, N and Sridhar, K.R. Seasonal and Geographical variations in the antimicrobial activity of seaweeds from the Mangalore coast in India. *Bot. Mar;* 1991; 34: 279-284.
- [33] Cody V; Middleton E; Harborne JB; Baretze A. Plant flavonoids. In: *Biology and medicine II: Biochemical, cellular and medicinal properties.* Alan R. Liss Inc, New York, 1988, 330.
- [34] Nyarko AA, Addy ME. Effects of aqueous extract of *Adenia cissampeloides* on blood pressure and serum analyse of hypertensive patients. *Phytotherapy Res*, 1990; 4 (1): 25-28.
- [35] Shimada T. Salivary proteins as a defense against dietary tannins. *J. Chem. Ecol.*, 2006; 36(6): 1149-1163.
- [36] Ketzis JK; Vercruyse J; Stromberg BE; Larsen M; Athanasiadou S; Houdijk JG. Evaluation of efficacy expectations for novel and non-chemical helminth control strategies in ruminants. *Vet. Parasitol.*, 2006; 139, 321-335.
- [37] Koleckar V; Kubikova K; Rehakova Z; Kuca K; Jun D; Jahodar L; Opletal L. Condensed and hydrolysable tannins as antioxidants influencing the health. *Mini Rev. Med. Chem.*, 2008; 8: 436-447.
- [38] Buzzini P, Arapitsas P, Goretti M., Branda E., Turchetti B., Pinelli P, Ieri F, and Romani A. Antimicrobial and antiviral activity of hydrolysable tannins, *Mini. Rev. Med. Chem.*, 2008; 8: 1179-1187.
- [39] Chung K.T., Wong T.Y., Wei C.I., Huang Y.W., Lin Y., Tannins and human health: A review. *Crit. Rev. Food Sci. Nutr.* 1998; 38:421–464.

- [40] Chakkaravarthy and Kumar. HPTLC and finger print analysis of steroid, flavonoid, and anti radical activity in *Sargassum wightii* for Gulf of Mannar, *Food Chemistry*, 2009; 124, 1147-1732.
- [41] Omulokoli E, Khan B, Chhabra SC. Antiplasmodial activity of four Kenyan medicinal plants. *J. Ethnopharmacol.* 1997; 56: 133-137.
- [42] Cowan, M.M. Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 1999; 12: 564-582.
- [43] Srivastava, N., K. Saurav, V. Mohanasrinivasan, K. Kannabiran and M. Singh., (2010). Antibacterial Potential of Macroalgae Collected from the Mandapam Coast, *India British J. Pharmacol. and Toxicol.*, 1(2): 72-76.
- [44] Nobori T; Miurak K; Wu DJ; Takabayashik CA; Carson DA. Deletions of the cyclin-dependent kinase-4 inhibitor gene in multiple human cancers *Nature*, 1994, 46, 753-756.
- [45] Stray F. The natural guide to medicinal herbs and plants. Tiger Books International, London, 1998:12-16.
- [46] DE Okwu and ME Okwu. Phytochemicals and vitamin content of indigenous spices of South Eastern Nigeria. *J. Sustain. Agric. Environ.*, 2004; 6(2): 140-147.
- [47] Sanchez-Machado DI; Lopez-Hernandez J; Pa Seiro Losada P; J Lopez-Cervantes. An HPLC method for the quantification of sterols in edible seaweeds *Biomed. Chromatogr.*, 2004; 18: 183-190.
- [48] Syakirah Zulkifi K; Abdullah N; Abdullah A; Aziman N; Wan Kumarudin W. Phytochemical Screening and activities of hydrophilic and lipophilic antioxidant of some fruit peels, *The Malaysian journal of Analytical Science* 2012, 16(3), 309-317.
- [49] Bhuvaneswari, S and Murugesan, S. In vitro antioxidant activity of marine red algae *Chondrococcus hornemanni* and *Spyridia fusiformis*; *Journal of Chemical and Pharmaceutical Research*, 2013; 5(3):82-85.
- [50] Manchu N., Melpha Y. and Edwin James J. Phytochemical investigation of three species of *Ulva* from Rasthacaud Coast, Tamil Nadu, India. *Journal of Chemical and Pharmaceutical Research*, 2014; 6(8):570-574.
- [51] Kuda, T and Ikemori, T. Minerals, polysaccharides and antioxidant properties of aqueous solutions obtained from macro algal beach casts in the Noto Peninsula, Ishikawa, Japan. *Food. Chem.*, 2009; (112): 575-581.