

Research Article

In vitro antibacterial effects of red alga *Champia parvula* (C. Agardh) of various solvents against human pathogenic bacteria

Vinoth Kumar, R; Murugesan, S^{*}; Bhuvaneshwari, S; Thennarasan, S

Unit of Algal Biotechnology and Bionano Technology, PG and Research Dept of Botany, Pachaiyappa's College, Chennai – 600 030

***Correspondence Info:**

Murugesan, S,
Unit of Algal Biotechnology and
Bionano Technology, PG and
Research Department of Botany,
Pachaiyappa's College, Chennai –
600 030
Email: smurugesan5@gmail.com

Keywords:

Champia parvula,
Methanol extract,
Pathogenic bacteria,
Antimicrobial activity

Abstract

The aim of the present study is to evaluate the antimicrobial inhibitory effect of *Champia parvula* (red alga) of various solvents at the concentration 100 µg/mL, on pathogenic bacteria like *Klebsiella pneumoniae*, *Proteus vulgaris*, *Bacillus cerus*, *Bacillus subtilius*, *Staphylococcus aureus*, and *Salmonella typhii* were studied by the disc diffusion method. The present study reveals that a higher zone of inhibition against *Salmonella typhii* (15.4 ± 0.2), *Bacillus subtilius* (13.8 ± 0.1), *Staphylococcus aureus* (10.7 ± 0.2) and *Proteus vulgaris* (10.6 ± 0.1) in the methanol extract alone, followed by acetone, benzene, chloroform, and ethyl acetate extracts showed moderate activity against most of the pathogens, whereas chloroform extract is inactive only against *Bacillus cerus*. The positive control streptomycin shows inhibitory action against all the pathogens studied. This study shows the potential of marine active compounds from *Champia parvula* as an antimicrobial agent for a disease free environment.

1. Introduction

Bacteria are the causative agents for many diseases among human population. Multi-drug resistant bacteria have evolved during recent times and hence, the searches for new drugs from natural resources have been in search for combating the diseases. Marine algae are the major resource available from the marine ecosystem and contain various biologically active compounds which serve as a component for nutraceutical and pharmaceutical industry [1]. Now, more than 2400 marine natural products have been isolated from seaweeds of subtropical and tropical populations [2]. Marine macroalgae are considered to be an excellent source of bioactive compounds with broad range of biological activities including antibacterial [3][4], antifungal [5], antiviral [4][6], antitumor [7], antioxidant [8][9][10], and anti-inflammatory activities [11][12][13]. Marine macroalgae and its active constituents such as carotenoids, terpenoids, steroids, amino acids, phlorotannins, phenolic compounds, halogenated ketones, alkanes and cyclic polysulphide find their applications in pharmaceutical, food, cosmetic industries and in traditional medicine [14][15][16]. Many bioactive compounds of marine algae with antimicrobial activity have been isolated and some of them are under investigation to protect lifestyle related diseases, [6][14][17][18][19]. Seaweeds are considered to produce a great variety of secondary metabolites characterized by a broad spectrum of biological activities. The compounds with cytostatic, antiviral, antihelminthic, antifungal and antibacterial activities have been detected in green, brown and red algae [20][21]. Extracts of marine algae were reported to exhibit antibacterial activity [22][23][24]. Several workers have reported that the seaweed extracts exhibit inhibitory activity against a number of gram positive and gram negative bacterial pathogens. A number of seaweeds have been studied for their antibacterial activity both in India and abroad [25][26][27][28][29][30]. India has a long coastline and abundant natural resources of marine algae with very high species diversity. The aim of this study was to analyse the antibacterial efficacy using acetone, benzene, chloroform, methanol, and ethyl acetate crude extracts of marine red alga *Champia parvula* (Rhodophyta) from the Mandapam coast of Tamil Nadu, against bacterial pathogens.

2. Materials and Methods

2.1 Collection of seaweed

The marine red alga *Champia parvula* (C. Agardh) was collected from Mandapam coast, Ramanathapuram District, and Tamil Nadu, India during low tides. The healthy epiphytic free submerged thalli were collected and the debris such as sand particles, pebbles and shells were removed. The seaweed samples were cleaned with sea water and then with fresh sea water and shade dried at room temperature. The dried samples were thoroughly washed with sterile sea water, followed by a rapid rinse with distilled water to remove the salt on the surface of the thalli. Excess water was removed with blotting paper, dried at room temperature until constant weight obtained and ground in an electric mixer and were used for the experimental procedures. The sample was identified by Emeritus Professor, Dr. M. Baluswamy, Associate Professor in Plant Biology and Plant Biotechnology, Madras Christian College, University of Madras, Chennai.

2.2 Preparation of extracts

The powdered sample (100g) was extracted in soxhlet apparatus using five solvents acetone, benzene, chloroform, methanol and ethyl acetate (1000 mL) as solvents for 24hrs at a room temperature maintained not more than the boiling point of the solvent. The resultant crude extracts were filtered with Whatman No.1 filter paper. The filtrates obtained were concentrated under vacuum with a rotary evaporator to obtain the various crude extracts. The crude extracts were collected in an air tight container and stored at 4°C until use.

2.3 Test Microorganisms

The strains of *Klebsiella pneumoniae*, *Proteus vulgaris*, *Bacillus cereus*, *Bacillus subtilis*, *Staphylococcus aureus*, and *Salmonella typhi* were obtained from the University of Madras, Taramani campus, Chennai, Tamil Nadu, India, were used as the test organisms for the study.

2.4 Preparation of Inoculum

The Nutrient agar slant slopes stored 4°C were used for inoculum preparation and antibacterial assay activity. The virulent bacterial cultures used for the assay were prepared by transferring a loopful of cells from the stock cultures. The bacteria were sub cultured and routinely maintained on both nutrient agar and Muller-Hinton agar. The antibacterial activity was carried out by was used. Antimicrobial activity was evaluated using Muller-Hinton agar by Kirby-bauer method. The various extracts 100 µg/mL was loaded on sterile filter paper discs and air dried. The bacterial pathogens were streaked on Muller-Hinton agar plates with sterile loops, the discs were then placed on seeded medium plates and incubated for 24 hours at 37°C for clear zone of inhibition. Sterile commercial disc were used as control. [12].

2.5 Antibacterial assay

The antibacterial activity of the algal extract was evaluated by Disc diffusion method[31]. A sterile cotton swab was dipped into the bacterial suspension and evenly streaked over the entire surface of a sterile Muller Hinton agar plate to obtain uniform inoculums. Sterile disc (6mm in diameter) loaded with various crude extract (100 µg/ml) were impregnated on to the plates and incubated overnight at 37°C for 24 hrs. The antibacterial activity was determined by measuring the diameter of the zone of inhibition (mm).

3. Results and Discussion

The antibacterial activity of various solvent extracts of *Champia parvula* (C. Agardh) on bacterial pathogens were studied by the Kirby-Bauer method and the results were measured in Plate-1 and tabulated in Table-1. The maximum inhibitory activity was observed with methanolic extract followed by other solvent extracts at the concentration 100 µg/ml. Streptomycin was used as the positive control in the study. The antibacterial activity of *Champia parvula* methanolic extract against bacterial pathogens are: *Salmonella typhi* (15.4 ± 0.2 mm), *Bacillus subtilis* (13.8 ± 0.1 mm), *Staphylococcus aureus* showed (10.7 ± 0.2 mm), and *Proteus vulgaris* (10.6 ± 0.1 mm) followed by other solvents such as acetone, benzene, chloroform and ethyl acetate which showed minimal antibacterial activity. No activity was observed with chloroform extract for the *Bacillus cereus*. The positive control streptomycin showed resistance to all the bacterial pathogens used in the study.

The marine red algae constituted active compounds which showed antibacterial activity [32][33]. Infections have become the leading cause of death worldwide which has led to an increase in antibacterial resistance, making it a global growing problem nowadays. Thus, there is a need to discover new antimicrobial compounds from natural resources with diverse bioactive compounds and novel mechanisms of action for new and emerging infectious diseases [34]. Reports have

suggested that the samples of powdered fresh or dried seaweeds showed less antibacterial activity than the extracts of powdered samples.

Bioactive and pharmacologically important compounds such as alginate, carrageen and agar as phycocolloids have been obtained from seaweeds and used in medicine and pharmacy have been reported many researchers [22]. Antibacterial activity of the extracts of marine red algae have been reported by other authors [35][36][37]. The results of the present study were direct evidence of *Champia parvula* possessing antibacterial activity which may be due to the presence secondary metabolites and phytochemicals present in the extracts.





Plate.1 Antibacterial activity of the different solvents extracts of *Champia parvula*

= A- Acetone, B- Benzene, Ch-Chloroform, M-Methanol, Ea- Ethyl acetate, S-Streptomycin
 = Kp - *Klebsiella pneumoniae*; Pv- *Proteus vulgaris*; Bc- *Bacillus cereus*; Bs- *Bacillus subtilis*;
 Sa- *Staphylococcus aureus*; St- *Salmonella typhi*
 = Cp-*Champia parvula*

Table 1: Antibacterial activity of various solvent extracts of *Champia parvula* against bacterial pathogens

Solvents	Bacterial pathogens showing zone of inhibition (mm in diameter)					
	<i>Klebsiella pneumoniae</i>	<i>Proteus vulgaris</i>	<i>Bacillus cereus</i>	<i>Bacillus subtilis</i>	<i>Staphylococcus aureus</i>	<i>Salmonella typhi</i>
Acetone	5.3 ± 0.0	5.4 ± 0.2	6.2 ± 0.1	5.6 ± 0.2	5.4 ± 0.4	6.5 ± 0.2
Benzene	5.4 ± 0.2	6.5 ± 0.2	7.4 ± 0.2	7.4 ± 0.2	8.3 ± 0.3	6.2 ± 0.0
Chloroform	7.4 ± 0.1	9.3 ± 0.1	0.0 ± 0.0	10.6 ± 0.1	9.6 ± 0.2	6.3 ± 0.2
Methanol	7.2 ± 0.0	10.6 ± 0.1	6.5 ± 0.2	13.8 ± 0.1	10.7 ± 0.2	15.4 ± 0.2
Ethyl acetate	7.5 ± 0.2	10.6 ± 0.1	6.5 ± 0.2	13.3 ± 0.2	5.5 ± 0.3	7.3 ± 0.2
Positive control Streptomycin	16.3 ± 0.1	9.2 ± 0.1	7.7 ± 0.1	11.7 ± 0.5	18.7 ± 0.2	19.2 ± 0.8

- = No activity, mm = millimeter, values, are expressed as mean ± standard error (n=3)



4. Conclusion

From the above results, it can be concluded that the marine red alga *Champia parvula* (C. Agardh) shows maximum activity to the bacterial pathogens with methanolic extract compared to other solvents which shows its potential as a good alternative source as antibacterial agents among red seaweeds from the natural source.

References

- [1] Kandhasamy, M. and K.D. Arunachalam. Evaluation of in vitro antibacterial property of seaweeds of southeast coast of India. *African J. Biotech.*, 2008; 7(12): 1958-1961.
- [2] Manilal A, and Sujith S. Biopotentials of Seaweeds Collected From Southwest Coast Of India. *J Mar Sci Tech.* 2009; 17: 67-73.
- [3] Singh, A. and Chaudhary, B. Preliminary phyco-chemical analysis and in vitro antibacterial screening of *Pithophora oedogonia* (Mont.) Wittrock: A freshwater green alga forming mats in the water bodies. *Journal of Algal Biomass Utilization*, 2010; 1: 33-41.
- [4] Bouhlal, R., Riadi, H. and Bourgougnon, N. Anti- viral activity of the extracts of Rhodophyceae from Morocco. *African Journal of Biotechnology* 2010; 9: 7968-7975.
- [5] De Felício, R., Dealbuquerque, S., Young, M.C.M., Yo-koya, N.S. and Deboni, H.M. Trypanocidal, leishmanicidal and antifungal potential from marine red alga *Bostrychia tenella* J. Agardh (Rhodomelaceae, Ceramiales). *Journal of Pharmaceutical and Biomedical Analysis*, 2010; 52: 763-769
- [6] Bouhlal, R., Haslin, C., Chermann, J.C., Collic-Jouault, S., Siquin, C., Simon, G., Cerantola, S., Riadi, H. and Bourgougnon, N. Antiviral activities of sulfated polysaccharides isolated from *Sphaerococcus coronopifolius* (Rhodophyta, Gigartinales) and *Boergeseniella thuyoides* (Rhodophyta, Ceramiales). *Marine Drugs*, 2011; 9: 1187-1209.
- [7] Kim, S.K. and Karadeniz, F. Anti-HIV activity of extracts and compounds from marine algae. *In Advanced Food and Nutrition Research*, 2011; 64: 255-265.
- [8] Kudaa, T., Tsunekawaa, M., Gotoa, H. and Arakib, Y. Antioxidant properties of four edible algae harvested in the Noto Peninsula, Japan. *Journal of Food Composition and Analysis*, 2005; 18: 625-633.
- [9] Alghazeer, R., Whida, F., Al-Najjar, A., Majdoob, H. and Al-Mazoghi, E. Assessment of antioxidant activity and phenolic content of some marine algae from the Libyan Coast. *Ain Shams Science Bulletin*, 2008; 46: 67-78.
- [10] Devi, G.K., Manivannan, K., Thirumaran, G., Rajathi, F.A.A. and Anantharaman, P. *In vitro* antioxidant activities of selected seaweeds from southeast coast of India. *Asian Pacific Journal of Tropical Medicine*, 2011; 4: 205- 211.
- [11] Vineela, C.H. and Elizabeth, K.M. Antimicrobial activity of marine algae of Visakhapatnam city, Andhra Pradesh. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences*, 2005; 7: 209-212.

- [12] Inci Tuney, Bilge Hilal Cadirci, Dilek Unal, Atakan Sukatar. Antimicrobial activities of the extracts of marine algae from the coast of Urla (Izmir, Turkey). *Turk J Biol*, 2006; 30: 171-175.
- [13] Patra J.K., Rath S.K., Jena, K., Rathod, V.K. and Thatoi, H. Evaluation of antioxidant and antimicrobial activity of seaweed (*Sargassum* spp.). Extract: A study on inhibition of glutathione-S-transferase activity. *Turkish Journal of Biology*, 2008; 32: 119-125.
- [14] Taskin, E., Ozturk, M., Taskin, E. and Kurt, O. Antibacterial activities of some marine algae from the Aegean Sea (Turkey). *African Journal of Biotechnology*, 2007; 6: 2746-2751.
- [15] Guedes, A., Amaro, M. and Malcata, F. Microalgae as Sources of Carotenoids. *Marine Drugs*, 2011; 9: 625-644. doi:10.3390/md9040625.
- [16] Fitton, J.H. Antiviral properties of marine algae. In: Critchley, A.T., Ohno, M. and Largo, D.B., Eds., *World Seaweed Resources*, ETI Information Services, Woking-ham, 2006: 7.
- [17] Mtolera, M.S.P. and Semesi, A.K. Antimicrobial activity of extracts from six green algae from Tanzania. In: Björk, M., Semesi, A.K., Pedersén, M. and Bergman, B., Eds., *Current Trends in Marine Botanical Research in the East African Region*, SIDA 1996; 211-217.
- [18] Bhacuni, D.S. and Rawat, D.S. Bioactive marine natural products. Springer, New York, 400. *Biology*, 1997; 36: 638-643
- [19] Priyadarshini, S., Bragadeeswaran, S., Prabhu, K. and Ran, S.S. Antimicrobial and hemolytic activity of seaweed extracts *Ulva fasciata* (Delile 1813) from Mandapam, Southeast coast of India. *Asian Pacific Journal of Tropical Biomedicine*, 2011; 1: S38-S39.
- [20] Lindequist U., Schweder T. Marine Biotechnology. In: Rehm, H. J., Reed, G. (Eds.), *Biotechnology*, vol. 10. Wiley-VCH, Weinheim, pp. 2001: 441-484
- [21] Newman DJ., Cragg GM., and Snader KM. Natural products as source of new drugs over the period 1981-2002. *J. Nat. Prod.* 2003; 66:1022-1037.
- [22] Siddhanta AK, Mody KH, Ramavat BK, Chauhan VD, Gara HS, Goel AK, Doss MJ, Srivasta MN, Patnaik GK, Kamboj VP. Bioactivity of marine organisms: part VIII – Screening of some marine flora of western coast of India. *Indian Journal of Experimental Biology*, 1997; 36: 638-643.
- [23] Mahasneh I., Jamal M., Kashashneh, M., Zibdeh M. Antibiotic activity of marine algae against multiantibiotic resistant bacteria. *Microbios.* 1995; 83:23-26.
- [24] Sachithananthan K., and Sivapalan A. Antibacterial properties of some marine algae of Sri Lanka. *Bulletin of Fisheries Research Station*, Sri Lanka. 1975; 26:5-9.
- [25] Pratt R., Mautner R., Gardener GM., Sha Y., Dufrenoy J. Report on antibiotic activity of seaweed extracts. *J. Amer. Pharm. Asst. Sci.* 1951; 40:579-579.
- [26] Chester CG., Stott JA. The production of antibiotic substances by seaweeds. In: (T.Braarud and N.A. Soressen, eds) *Second International Seaweed Symp.* Pergamon, Newyork. 1956 pp.49-53
- [27] Burkholder PR., Burkholder LM., Almodovar LR. Antibiotic activity of some marine algae of Puerto Rico. *Bot. Mar.* 1960; 2: 149-156.
- [28] Allen MB., Dawson NY. Production of antibacterial substances by benthic tropical marine algae. *J.Bacteriol.* 1960; 79: 459-460.
- [29] Nunez G., and Sanabria LMS. Effects of products by extracts of macroscopic marine algae on bacterial growth. *Acta. Biol. Venez* 1975;. 9:121-134.
- [30] Sitakara Rao V., and Tipnis VK. Protein content of marine algae from Gujarat coast. **Curr. Sci.** 1964; 33: 16-17.
- [31] Bauer, A.W., Kirby, W.M., Sherris, J.C. and Turck, M. Antibiotic susceptibility testing by a standardized single disk method. *American Journal Clinical Pathology*, 1996; 45: 493-496.
- [32] Del Val, A.G., Platas, G., Basilio, A., Cabello, A., Gorro-chateui, J., Suay, I., Vicente, F., Portillo, E., DeRio, M.J., Reina, G.G. and Pelaez, F. Screening of antimicrobial activities in red, green and brown macroalgae from Gran Canaria (Canary Islands, Spain). *International Microbiology*, 2001; 4: 35-40.
- [33] Lavanya, R. and Veerappan, N. Antibacterial potential of six seaweeds collected from gulf of mannar of southeast coast of India. *Advances in Biological Research*, 2011; 5: 38-44.
- [34] Adaikalaraj G, Patric Raja D, Johnson M, Janakiraman N, Babu A. Antibacterial potential of selected red seaweeds from Manapad coastal areas, Thoothukudi, Tamil Nadu, India. *Asian Pacific Journal of Tropical Biomedicine.*

2012S1077-S1080.

- [35] Rajasulochana P, Dhamotharan R, Krishnamoorthy P, Murugesan S. Antibacterial Activity of the Extracts of Marine Red and Brown Algae. *J Amer Sci*; 2009; 5: 20-25.
- [36] Krishnapriya Madhu Varier, Muthu Chinnadurai John Milton, Chinnasamy Arulvasu and Babu Gajendran. Evaluation of antibacterial properties of selected red seaweeds from Rameshwaram, Tamil Nadu, India *J. Acad. Indus. Res.* 2013: Vol. 1(11).
- [37] Shanmuga Priya S, Krishnareni J, Joseph Selvin, Gandhimathi R, Arun Kumar M, Thangavelu T, Seghal kiran G, Natarajaseenivasan K). Optimization of extracellular thermotolerant alkaline protease produced by marine *Roseobacter* Sp (MMD040). *Bioprocess Biosyst Eng.*, 2008; 31: 427-433.