



## 5. Conclusion

ESR is increased in pregnancy because of elevated plasma globulins and fibrinogen. Hemodilution during pregnancy may be another cause of increased ESR.

## References

1. Pal G.K, Textbook of Medical Physiology 2<sup>nd</sup> edition. p491-92.
2. Litchman M. A. Beutler E, Kipps TJ, Seligsohn URI, Kaushanky K, Prchal JT. Williams Hematology. 7ed, USA McGraw Hill.2006.p.101.
3. Dutta.D.C Textbook of obstetrics.6<sup>th</sup>ed.kolkatta: New central book agency, 2004.p51-52.
4. Taylor.D.J, Lind,T. Puerpural Haematological indices. *Br J Obstet Gynaecol.* 1981; 88(6):601-606.
5. Manten GTR, Franx A, sikkema JM, et al. Fibrinogen and high molecular weight fibrinogen during and after normal pregnancy. *Thrombosis Res.* 2004, 114:19.
6. Hytten FE, Leitch I. The physiology of Human pregnancy 2<sup>nd</sup> ed. Philadelphia, Davis, 1971.
7. Johnson RL. Thromboembolic diseases complicating pregnancy. In Foley MR, Strong TH (eds).Obstretic Intensive care. A practical manual. Philadelphia, WB saunders company.1997:91
8. Ozanne p, Linderkamp o, Miller F, et al, Erythrocyte aggregation during normal pregnancy. *Am J Obstet Gynecol.*1983; 147:576.