

Development of validated HPTLC method for the estimation of betasitosterol in marketed herbal formulation of muscle and joint HRX pain relieving oil

M. S. Charde^{1*}, M. D. Chakolkar¹, A. S. Welankiwar¹, U. Keshwar² and B. K. Shrikande²

¹Government College of Pharmacy Kathora Naka, Amravati.444604 MS India

²Baidyanath Life Sciences Pvt. Ltd Nagpur

*Correspondence Info:

Dr. M. S. Charde

Government College of Pharmacy,

Kathora Naka, Amravati.444604 MS India

E-Mail: manojdps@rediffmail.com

Abstract


A new and simple HPTLC method was developed and validated for the quantitative estimation of betasitosterol in muscle and joint pain relaxant herbal oil. TLC aluminium plates precoated with silica gel 60F-254 (0.2 mm thickness) were used. The linear ascending development was carried out in twin trough glass chamber saturated with mobile phase Toluene: Ethyl acetate (7:3) ratio followed by densitometric determination was carried out by TLC scanner (CAMAG) at 560 nm in reflectance/absorbance mode. The R_f value was found to be 0.58. Linearity was found to be in the concentration range of 500 ng to 3000 ng. The linear regression data for the calibration plots showed a good linear relationship with r²=0.99 for Betasitosterol. According to the ICH guideline the method was validate for accuracy, precision, Specificity and ruggedness. The proposed method is accurate, precise, reproducible, and can be adopted for routine analysis of Betasitosterol from herbal joint and muscle pain relieving oil by HPTLC.

Keywords: Betasitosterol, HPTLC, Joint, Muscle, Herbal oil, Densitometric

1. Introduction

High Performance Thin Layer Chromatography (HPTLC) is a powerful method equally suitable for qualitative and quantitative analytical tasks. Applications of HPTLC, such as identification and quantitation of constituents, impurities, active substances, process development and optimization, process monitoring, and cleaning validation have been demonstrated. HPTLC has been reported to provide excellent separation, qualitative and quantitative analysis of a wide range of compounds, such as herbal and botanical dietary supplements, nutraceuticals, traditional western medicines, traditional Chinese medicines and Ayurvedic (Indian) medicines and determination of radio labeled substances in chemical, biochemical, biological, pharmaceutical, and medicinal samples. HPTLC is superior to other analytical techniques in terms of total cost and time for analysis. It is an offline process in which various stages are carried out independently. Important features of HPTLC include the ability to analyze crude samples containing multi-components, application of large number of sample and a series of standards using the spray-on technique, choice of solvents for the HPTLC development is wide as the mobile phases are fully evaporated before the detection step, processing of standards and samples identically on the same plate leading to better accuracy and precision of quantification, different and universal selective detection methods, and in situ spectra recording in sequence to obtain positive identification of fractions, storage of total sample on layer without time constraints¹⁻³. The drug undertaken⁴ for the study is Betasitosterol in Marketed formulation (HRX) from Baidyanath Life Sciences Pvt. Ltd Nagpur. Betasitosterol [Figure 1] Chemically is 17-(5-Ethyl-6-methylheptan-2-yl)-10,13-dimethyl-2,3,4,7,8,9,11,12,14,15,16,17- dodecahydro- 1Hcyclopenta [a] phenanthren-3-ol. And it is used in treatment of elevated blood cholesterol levels also used in prostatic hyperplasia. As the literature survey⁵⁻¹⁰ clearly reveals that there is no proper analytical method available for the quantitative estimation of betasitosterol in polyherbalmuscle and joint pain relieving oil, thus the present study aims to develop a rapid, efficient and reproducible method of analysis for betasitosterol in polyherbal muscle and joint pain relieving oil by HPTLC.

Figure 1: Chemical structure of Betsitosterol



2. Experimentals

2.1 Materials and Methods

Markers were used for analytical method development and quantitative analysis. They were purchased from supplier Natural Remedies Pvt Ltd., by Baidyanath Life Sciences Pvt. Ltd. Analytical grade of toluene, ethyl acetate, methanol, pet. Ether were used. Stationary phase was pre-coated silica gel aluminium plate 60 F₂₅₄ was obtained from Merck, Germany.

2.2 Instrumentation and Chromatographic Conditions

The chromatography was performed on (10 cm x 10 cm) aluminum plate coated with (0.2 mm) silica gel 60 F254 (E. Merck, Germany). The samples were applied to the plate as bands width 6 mm by using Camag (Muttentz, Switzerland) Linomat 5 applicator fitted with 100 µl syringe (Hamilton, Switzerland). The rate of application was constant at 150nL/s and space between two bands was 15 mm. The mobile phase was used as e toluene: ethyl acetate (7:3), v/v). The Linear ascending development of the plate was carried out in twin-trough glass chamber previously saturated with mobile phase for 20 min at room temperature (250 ± 2) and relative humidity (60 % ± 5). The length of chromatogram run was 80 mm. After development, the plate was removed and the plate was air dried and derivatives with Anisaldehyde-Sulphuric acid. The densitometric scanning was performed at 560 nm using Camag TLC.

2.3 Determination of Betsitosterol in Oil Formulation¹⁻³

Preparation of Reference Standard Solution: An accurately weighed quantity Betasitosterol (1 mg) was dissolved in methanol (10 mL) and was sonicated for 5 min. gave the conc. of 100 ng/µl.

Preparation of Test Solution of Oil Formulation: Accurately weighed oil (1.0 g) was dissolved in pet ether (10.0 mL) and was sonicated for 5min. The solution was filtered through Whatman filter paper no. 1. From the above solution 1mL of this solution taken the volume was made up to 10.0 mL with pet ether, the final conc. of soln.was 10 ug/ µl.

Selection of Mobile Phase: For the HPTLC method development mobile phase selection different solvent system have been tried on trial and error base on aluminum plate coated with (0.2 mm) silica gel 60 F254 (E. Merck, Germany). Finally from above mention system Tolune: Ethyl acetate in ratio 9.3:0.7 give satisfactory result which gives RF value 0.51.

Linearity and Range: From the stock solution (100 ng/ml of betasitosterol) aliquots of 5, 10, 15, 20, 25 and 30 µl were spotted on TLC plate under nitrogen stream using Linomat Applicator, to obtain final concentration range 500-3000 ng/band for betasitosterol.The plate was dried in air and then the plate was developed in twin through developing chamber (20*10 cm) with stainless lead. Previously saturated with the mobile phase with 30 min. the plate was removed from chamber and dried in air and was scanned and quantified at 560 nm in excitation mode with camag scanner having software cats4. The peak areas were measured for each lane at Rf 0.58 ± 0.02 for betasitosterol. And the calibration curve of peak area VS concentration was plotted.

Estimation of Betsitosterol in Marketed Formulation: The content of Betasitosterol was determined by applying an appropriate volume of 10 µL and 10 µL of test solution and standard solution, respectively. The plate was developed and scanned as per proposed chromatographic conditions. The conc. was determined by linear regression equation.

2.4 Method Validation¹⁷:

Accuracy: Accuracy of the proposed method was ascertained on the basis of recovery studies performed by standard addition method (spiking). Accurately measured amount of standard Betasitosterol was added on the sample tracks on TLC plate in subsequently increasing conc. of 500 ng/spot in Duplicate. The chromatogram was developed and scanned as per proposed chromatographic conditions. The percentage recovery of standard betasitosterol from the proposed method was calculated.

Precision:

Intra-day: Intraday precision was determined by analysis of betasitosterol standard solutions in the range 1000-3000 ng/band for three times on the same day and % RSD was calculated.

Inter-day: Interday precision was assessed by analyzing of the same solution on three different days over a period of one week and %RSD was calculated.

Repeatability: The repeatability of sample application was assessed by spotting 5 µL contain 500 ng/spot of standard Betsitosterol on TLC plate (n=6).The plate was developed and scanned as per proposed chromatographic conditions. The average, standard deviation (S.D.) and percentage relative standard deviation (% RSD) of peak area was calculated.

Specificity: The specificity of the method was studied by analyzing standards and formulation by simultaneously applying on the same plate. The spots of Betasitosterol in formulation were confirmed by comparing Rf values with that of reference standard. The peak purity of individual standard in sample track was assessed by comparing spectra at peak start, peak apex and Peak end positions of the spot.The chromatograms of formulation, reference standard and their overlain spectrum are shown, respectively in figure 4,5,6.

Ruggedness: The ruggedness of method was done at conc. levels 1500 ng/spot of working standard solution of Betasitosterol The values of % RSD was lower than 2 indicate the ruggedness of the method.

3. Results and Discussion

3.1 Determination of Betasitosterol in Oil Formulation:

Selection of Mobile Phase

Table 1: Mobile phase selection data for betasitosterol

Sr No.	Name of solvent system	Ratio	Rf
1	Toluene: Ethyl acetate	7:3,9:1	0.51
2	Toluene: Methanol	9:1	-
3	Tolune:Methanol: pet ether	6:3:1	-

Figure 2: Densitometric scan of Betsitosterol at 560 nm.




Figure 3: Linearity study of betasitosterol




Table 2: Linearity study of betasitosterol

Conc. in ng	500	1000	1500	2000	2500	3000
Area	2754.6	3345.4	3956.8	4562.1	5126.6	5765.5

Estimation of Betasitosterol in Marketed Formulation

Table 3: Results of Marketed formulation analysis

Sr. No	formulation (ug)	Marker (ng)	Area
1	100	0	1284.7
2	100	0	1288.7
3	100	0	1285.4
4	100	0	1286.5
5	100	0	1287.1
6	100	0	1288.3
7	100	0	1287.4
8	0	1000	3345.4
Mean			1286.871
Std			1.456839
Amount found(ng)			257.36
Amount found (%)			0.38

3.2 Method Validation

Accuracy:

Table 4: Results of Recovery studies

Sr. No	Sample (ug) [A]	Area [B]	Amount found [C]	Standard added (ug/spot) [D]	Total applied [E]	Area [F]	Total recovered [G]	%recovery [G/E*100]
1	100	1286	257.36	500	757.63	4040.6	756.02	99.7539
2	100	1286	257.36	500	757.63	4040.6	756.02	99.7539
3	100	1286	257.36	1000	1257.36	4631.4	1256.83	99.96236
4	100	1286	257.36	1000	1257.36	4631.4	1256.83	99.96236
5	100	1286	257.36	1500	1757.36	5240.8	1757.06	99.99392
6	100	1286	257.36	1500	1757.36	5240.8	1757.06	99.99392
Mean (% recovered)							99.9	
SD (% recovered)							0.11	
%RSD (% recovered)							1.101101	

Precision:

Intra-Day:

Table 5: Results of Intra-day precision studies

Concentration (ng/spot)	Peak area± SD	Amount found	%Amount found
1000	3345.4±11.6	998.89	99.89
2000	3956.88±17	1998.72	99.36
3000	4562.1±16.6	2998.98	99.99

Inter-Day:

Table 6: Results of Inter-day precision studies

Concentration (ng/spot)	Peak area± SD	Amount found	%Amount found
1000	3332±11.83	998.69	99.69
2000	3946.55± 16.26	1998.39	99.91
3000	4542.1± 17.68	3000.41	100.01

Repeatability:

Table 7: Results of repeatability studies

Sr. No	conc. (ng/spot)	Area
1	500	2753.3
2	500	2753.9
3	500	2754.6
4	500	2754.1
5	500	2754.0
6	500	2753.7
Mean		2753.933
SD		0.394405
%RSD		0.014322

Specificity:

Figure 4: Chromatogram of Betsitosterol in oil formulation




Figure 5: Chromatogram of Standard Betsitosterol





Figure 6: Overlain spectrum of betasitosterol, standard and oil formulation



Ruggedness:

Table 8: Ruggedness data for betasitosterol

Drug	Analyst	Peak area (150ng/spot) Mean ± SD	% RSD	Amount Found	%Amount Found
Betasitosterol	I	5201±4.12	0.082	1500.32	100.21
	II	5209.5±4.57	0.091	1510.94	101.29

4. Conclusion

The newly developed method was found to be simple, specific, precise, rapid, and reproducible and can be used for quantification of betasitosterol in routine quality control of joint and muscle pain relieving polyherbal oil. The linearity and reproducibility data of the drugs carried out by this method showed that no major interference is caused in the estimation of the drug. The method is very simple and rapid and nowhere involves complicated sample preparation and mobile phase preparation. Also the proposed method showed good specificity and selectivity.

References

- Rashmin P et al. HPTLC Method Development and Validation: Strategy to Minimize Methodological Failures, *Journal of Food and Drug Analysis* 2012. 20(4).794-804.
- Patel, R. B., Patel, M. R. and Patel, B. G. Experimental aspects and implementation of HPTLC. In “Highperformance thin layer chromatography (HPTLC)”. Srivastava, M. M. ed. Springer Berlin Heidelberg, New York, USA. 2011. pp. 41-54.
- Sherma, J. Modern thin layer chromatography. 2008; *J. AOAC Int.* 91: 1142-1144.
- <http://en.wikipedia.org/wiki/Betasitosterol>.
- Tatke P et al. Marker based standardisation of commercial formulations and ethanolic extract containing Betasitosterol and D-Glucoside using HPTLC, *IRAJP* 2010. 1(2).616-623.
- Tatke et al. Development and Validation of A Novel HPTLC Method For Simultaneous Estimation Of Betasitosterol d glucoside and Withaferin A, *Int j Pharm Pharm Sci* 2011. 3(2).227-230.
- Unnati M. Shah et al. Development and Validation of a Simple Isocratic HPLC Method for Simultaneous Estimation of Phytosterols in *Cissus quadrangularis*, *Ind J Pharm Sci* 2010. 72(6).753-758.
- Niyati S Acharya et al. Development of Pharmacognostical parameters and estimation of betasitosterol using HPTLC in roots of *gmelina arborea roxb*, *phcog j* 2012. 4(30).1-12.
- Dighe V et al. Quantitation of B-Sitosterol from *Celastrus Paniculatus* Willd. Using Validated High Performance Thin Layer Chromatography Method, *Int J Pharm & Bio Sci* 2011. 2(2).117-123.
- Niranjan A, Tewari S.K. Phtyochemical composition and antioxidant activity of *Desmodium gangetium*, *Natural Product Radiance* 2008. 7 (1).35-3932.
- WHO guidelines for assessing quality of herbal medicines with reference to contaminants and residues, WHO Library Cataloguing-in-Publication Data, ISBN 978 92 4 159444 8, 2007.
- BishtAnupam, Zaman K., Singh, Mamta, Gupta Richa, Singh Vinod. Pharmacognostical studies on *Oroxylum indicum* (Linn.), *Indian Journal of Natural Products and Resources* 2011. 2(4).472-478.

13. Saraswathi R, Upadyay Lokesh, Venkatakrishnan R, Meera R, Devi P. phytochemical investigation analgesic and anti-inflammatory activity of *abutilon indicum*, *International Journal of Pharmacy and Pharmaceutical Sciences* 2011.3(2). 1-10
14. He Weiping and Huang Baokang .chemistry and bioactivities of a medicinal Spice: *Foeniculumvulgare*, *Journal of Medicinal Plants Research* 2011. 5(16). 3595-3600.
15. Indian Pharmacopoeia, Government of India, Ministry of Health and Family Welfare, Delhi, India; Part III (2007); Page no - 2071
16. Quantitative analysis of Drugs in Pharmaceutical Formulations, third edition 1997, Dr.P.D. Sethi, published by CBS publishers & distributors, New Delhi.
17. International Conference on Harmonization (2005) ICH harmonised tripartite guideline Validation of analytical procedures: text and methodology Q2 (R1) ICH, Geneva, Nov (2005).